

Anti-Insulin Receptor (phospho Tyr1150) + IGF1 Receptor (phospho Tyr1135) Antibody (A16411)

\sim					
~	മ	∩ıtı	\sim	tıc	ons:
$\mathbf{\mathcal{C}}$	บบ	UIII	Ca	u	nio.

Name: Anti-Insulin Receptor (phospho Tyr1150) + IGF1 Receptor (phospho Tyr1135) Antibody

Description: Rabbit polyclonal antibody to Insulin Receptor (phospho Tyr1150) + IGF1 Receptor

(phospho Tyr1135).

Applications: WB

Recommended Dilutions: WB: 1:500-1:2,000

Reactivity: Human, Mouse, Rat

Immunogen: A synthetic phosphorylated peptide around Y1150 of human IGF1RINSR

(NP_001073285.1).

Sequence: TDYYR

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 80 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium

Azide.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.



Anti-Insulin Receptor (phospho Tyr1150) + IGF1 Receptor (phospho Tyr1135) Antibody (A16411)

Images:



Western blot analysis of extracts of 293 and HepG2 cells, using Anti-Insulin Receptor (phospho Tyr1150) + IGF1 Receptor (phospho Tyr1135) Antibody (A16411) at 1:1,000 dilution. 293 cells were treated by EGF (25ug/mL) for 30 minutes after serum-starvation overnight. HepG2 cells were treated by EGF (100ng/ml) for 30 minutes after serum-starvation overnight. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% BSA.