


## Anti-UNG Antibody (A11430)

### Specifications:


Name:	Anti-UNG Antibody
Description:	Rabbit polyclonal antibody to UNG.
Applications:	WB, IHC
Recommended Dilutions:	WB: 1:1,000-1:2,000, IHC: 1:50-1:100
Reactivity:	Human, Mouse
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-313 of human UNG (NP_550433.1).
Sequence:	MIGQKTLYSFFSPSPARKRHAPSPEPAVQGTGVAGVPEESGDAAAIPAKKAPAGQEEP GTPPSSPLSAEQLDRIQRNKAALLRLAARNVPVGFGEVWKKHLSGEFGKPYFIKLMG FVAEERKHYTVYPPPHQVFTWTQMCDIKDVKVVILGQDPYHGPNQAHGLCFVQRPVP PPPSLENIYKELSTDIEDFVHPGHGDLGWAQGGVLLLNAVLTVRAHQANSHKERGWE QFTDAVWSWLNQNSNGLVFLWGSYAQKKGSAIDRKRHHVLTQTAHPSPLSVYRGFFGC RHFSKTNELLQKSGKKPIDWKEL
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	37 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-UNG Antibody (A11430)


### Images:



Western blot analysis of extracts of various cell lines, using Anti-UNG Antibody (A11430) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 5s.



Immunohistochemistry analysis of paraffin-embedded human stomach using Anti-UNG Antibody (A11430) at a dilution of 1:100 (40x lens).



Immunohistochemistry analysis of paraffin-embedded mouse brain using Anti-UNG Antibody (A11430) at a dilution of 1:100 (40x lens).