


Anti-ATG7 Antibody (A12684)

Specifications:


Name:	Anti-ATG7 Antibody
Description:	Rabbit polyclonal antibody to ATG7.
Applications:	IHC, ICC/IF
Recommended Dilutions:	IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 500-676 of human ATG7 (NP_006386.1).
Sequence:	LGFDTFVVMRHLKPKQGGAGDLCPNHPVASADLLGSSLFANIPGYKLGICYFCNDVV APGDSTRDRTLDDQCTVSRPGLAVIAGALAVELMVSVLQHPEGGYAIASSDDRMNEP PTSLGLVPHQIRGFLSRFDNVLPVSLAFDKCTACSSKVLQYEREGFNFLAKVFNSSH SFL
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	78 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-ATG7 Antibody (A12684)


Images:




Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using Anti-ATG7 Antibody (A12684) at a dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human colon tissue using Anti-ATG7 Antibody (A12684) at a dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse lung using Anti-ATG7 Antibody (A12684) at a dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat ovary using Anti-ATG7 Antibody (A12684) at a dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.