


## Anti-TRAF2 Antibody (A12723)

### Specifications:


Name:	Anti-TRAF2 Antibody
Description:	Rabbit polyclonal antibody to TRAF2.
Applications:	WB, IHC, IP
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:200, IP: 1:50-1:100
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human TRAF2 (NP_066961.2).
Sequence:	MAAASVTPPGSLELLQPGFSKTLGKLEAKYLCSACRNVLRRPFQAQCGHRYCSFCL ASILSSGPQNCAACVHEGIYEEGISILESSAFPDAARREVESLPAVCPDGDGCTWKG TLKEYESCHEGRCPMLTECPACKGLVRLGKERHLEHECPERSLSRHCRAPECCGAD VKAHHEVCPKFPLTCDGCGKKKIPREKFQDHVKTCGKCRVPCRFAIGCLETVEGEKQ QEHEVQWLREHLAMLLSSVLEAKPLLGDQSHAGSELLQRCESELEKKTATFENIVCVLN REVERVAMTA
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	54 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-TRAF2 Antibody (A12723)


### Images:




Western blot analysis of extracts of various cell lines, using Anti-TRAF2 Antibody (A12723) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST.



Immunohistochemistry analysis of paraffin-embedded mouse brain using Anti-TRAF2 Antibody (A12723) at a dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse spleen using Anti-TRAF2 Antibody (A12723) at a dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat spleen using Anti-TRAF2 Antibody (A12723) at a dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.