

Anti-G-protein coupled receptor 30 Antibody (A12843)

Specifications:

Name: Anti-G-protein coupled receptor 30 Antibody

Description: Rabbit polyclonal antibody to G-protein coupled receptor 30.

Applications: WB, ICC/IF

Recommended Dilutions: WB: 1:500-1:1,000, ICC/IF: 1:50-1:200

Reactivity: Human, Mouse, Rat

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 307-375

of human GPER1 (Q99527).

Sequence: HIVNLAAFSNSCLNPLIYSFLGETFRDKLRLYIEQKTNLPALNRFCHAALKAVIPDST

EQSDVRFSSAV

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 52 kDa


Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.

Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.

Images:




Western blot analysis of extracts of Mouse heart cells, using Anti-G-protein coupled receptor 30 Antibody (A12843) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 5s.




Anti-G-protein coupled receptor 30 Antibody (A12843)


Images continued:



Immunofluorescence analysis of NIH/3T3 cells using Anti-G-protein coupled receptor 30 Antibody (A12843) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of U20S cells using Anti-G-protein coupled receptor 30 Antibody (A12843) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of C6 cells using Anti-G-protein coupled receptor 30 Antibody (A12843) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).