

## Anti-Cathepsin B Antibody (A121151)

### Specifications:

Name:	Anti-Cathepsin B Antibody
Description:	Goat polyclonal antibody to Cathepsin B.
Applications:	ELISA, WB, IF, FC
Reactivity:	Human
Immunogen:	Synthetic peptide corresponding to Human Cathepsin B.
Sequence:	SDRICIHTNAHVSVE
Host:	Goat
Clonality:	Polyclonal
lsotype:	lgG
Conjugate:	Unconjugated
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Concentration:	100 μg at 0.5 mg/ml.
Product Form:	Liquid
Formulation:	Supplied in Tris Buffered Saline, 0.02% sodium azide, pH 7.3 with 0.5% BSA.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

### Images:


250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa 15kDa

Anti-Cathepsin B Antibody (A121142) (1 $\mu$ g/ml) staining of HepG2 cell lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.


# antibodies

## Anti-Cathepsin B Antibody (A121151)


#### Images continued:



Anti-Cathepsin B Antibody (A121142) (1µg/ml) staining of Human Lung (A) and HCORC (B) and (0.5µg/ml) HCBGLC (C) lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



Anti-Cathepsin B Antibody (A121142) - Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10µg/ml) followed by Alexa Fluor 488 secondary antibody (2µg/ml), showing cytoplasmic/vesicle and extracellular staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10µg/ml) followed by Alexa Fluor 488 secondary antibody (2µg/ml).



→ Anti- Cathepsin B

Anti-Cathepsin B Antibody (A121142) - Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10µg/ml) followed by Alexa Fluor 488 secondary antibody (1µg/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.