


## Anti-Adenosine Receptor A2a Antibody (A13527)

### Specifications:

Name:	Anti-Adenosine Receptor A2a Antibody
Description:	Rabbit polyclonal antibody to Adenosine Receptor A2a.
Applications:	WB, ICC/IF
Recommended Dilutions:	WB: 1:500-1:1,000, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human ADORA2A (NP_000666.2).
Sequence:	IDRYIAIRIPLRYNGLVTGTRAKGIIAICWVLSFAIGLTPMLGWNNCGQPKEGKNHSQ GCGEGQVACLFEDVVPMMNYMVYFNFFACVLVPLLLMLGVYLR
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	43 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


### Images:




Western blot analysis of extracts of SH-SY5Y cells, using Anti-Adenosine Receptor A2a Antibody (A13527) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.

## Anti-Adenosine Receptor A2a Antibody (A13527)


Images continued:



Immunofluorescence analysis of C6 cells using Anti-Adenosine Receptor A2a Antibody (A13527) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of HeLa cells using Anti-Adenosine Receptor A2a Antibody (A13527) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of NIH/3T3 cells using Anti-Adenosine Receptor A2a Antibody (A13527) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).