


Anti-GRK2 Antibody (A13580)

Specifications:


Name:	Anti-GRK2 Antibody
Description:	Rabbit polyclonal antibody to GRK2.
Applications:	WB, IHC
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 440-689 of human GRK2 (NP_001610.2).
Sequence:	LGRGAQEVKESPFRRSLDWQMVFLQKYPPPLIPPRGEVNAADAFDIGSFDEEDTKGIK LLDSDQELYRNFPLTISERWQQEVAETVFDTINAETDRLEARKKAKNKQLGHEEDYAL GKDCIMHGYMSKMGNPFLTQWQRRYFYLFPNRLEWRGEGEAPQSLTMEEQSVEETQ IKERKCLLLKIRGGKQFILQCDSPELVQWKELRDYREAQQLVQRVPMKMKPRSP VVLSKVPLVQRGSANGL
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	80 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-GRK2 Antibody (A13580)


Images:




Western blot analysis of extracts of various cell lines, using Anti-GRK2 Antibody (A13580) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST.



Western blot analysis of extracts from normal (control) and GRK2 knockout (KO) 293T cells, using Anti-GRK2 Antibody (A13580) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Immunohistochemistry analysis of paraffin-embedded rat brain using Anti-GRK2 Antibody (A13580) at a dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human esophageal cancer using Anti-GRK2 Antibody (A13580) at a dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.