

Anti-Factor XII Antibody (A13597)


Specifications:

| Name: | Anti-Factor XII Antibody |
|---|--|
| Description: | Rabbit polyclonal antibody to Factor XII. |
| Applications: | WB |
| Recommended Dilutions: | WB: 1:500-1:2,000 |
| Reactivity: | Human, Mouse |
| Immunogen: | Recombinant fusion protein containing a sequence corresponding to amino acids 336-615 of human F12 (NP_000496.2). |
| Sequence: | QTPGALPAKREQPPSLTRNGPLSCGQRLRKSLSSMTRVVGGLVALRGAHPYIAALYWG HSFCAGSLIAPCWVLTAAHCLQDRPAPEDLTVVLGQERRNHSCEPCQTLAVRSYRLHE AFSPVSYQHDLALLRLQEDADGSCALLSPYVQPVCLPSGAARPSETTLCQVAGWGHQF EGAEEYASFLQEAQVPFLSLERCSAPDVHGSSILPGMLCAGFLEGGTDACQGDSGGPL VCEDQAAERRLTLQGIISWGSGCGDRNKPGVYTDVAYYLAWIREHTVS |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Isotype: | lgG |
| Conjugate: | |
| Conjugate. | Unconjugated |
| Purification: | Unconjugated Affinity purification. |
| | |
| Purification: | Affinity purification. |
| Purification: Molecular Weight: | Affinity purification. 71 kDa |
| Purification: Molecular Weight: Product Form: | Affinity purification. 71 kDa Liquid Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium |

antibodies

Anti-Factor XII Antibody (A13597)

Images:



Western blot analysis of extracts of various cell lines, using Anti-Factor XII Antibody (A13597) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST.