


## Anti-Glutamate Receptor 1 (AMPA subtype) Antibody (A13683)

### Specifications:


Name:	Anti-Glutamate Receptor 1 (AMPA subtype) Antibody
Description:	Rabbit polyclonal antibody to Glutamate Receptor 1 (AMPA subtype).
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 330-470 of human GluR1/GRIA1 (NP_000818.2).
Sequence:	PWGQGIDIQRALQQVRFEGLTGNVQFNEKGRRRTNYTLHVIEMKHDGIRKIGYWNEDDK FVPAATDAQAGGDNSSVQNRTYIVTTILEDPYVMLKKNANQFEGNDRYEGYCVELAAE IAKHVGYSYRLEIVSDGKYGARDPD
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	100 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-Glutamate Receptor 1 (AMPA subtype) Antibody (A13683)


### Images:




Western blot analysis of various lysates, using Anti-Glutamate Receptor 1 (AMPA subtype) Antibody (A13683) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded mouse brain using Anti-Glutamate Receptor 1 (AMPA subtype) Antibody (A13683) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of rat brain cells using Anti-Glutamate Receptor 1 (AMPA subtype) Antibody (A13683) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of mouse brain cells using Anti-Glutamate Receptor 1 (AMPA subtype) Antibody (A13683) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).