


Anti-Syk Antibody (A13889)

Specifications:

Name:	Anti-Syk Antibody
Description:	Rabbit polyclonal antibody to Syk.
Applications:	WB, ICC/IF
Recommended Dilutions:	WB: 1:500-1:2,000, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 250-350 of human Syk (NP_001128524.1).
Sequence:	GLLRVLTVPCQKIGTQGNVNFGGRPQLPGSHPASSPAQGNRQESTVSFNPYEP LAPW AADKGPQREALPMDTEVYESPYADPEEIRPKEVYLDRKLLTLE
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	69 kDa / 72 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Western blot analysis of extracts of Rat liver, using Anti-Syk Antibody (A13889) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.

Anti-Syk Antibody (A13889)

Images continued:



Western blot analysis of extracts of various cell lines, using Anti-Syk Antibody (A13889) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 90s.



Immunofluorescence analysis of U2OS using Anti-Syk Antibody (A13889) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).