


Anti-Axin 2 Antibody (A14022)

Specifications:


Name:	Anti-Axin 2 Antibody
Description:	Rabbit polyclonal antibody to Axin 2.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:100, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 9-600 of human AXIN2 (NP_004646.3).
Sequence:	CLPDPSSSFREDAPRPPVPGEEGETPPCQPGVKGQVTKPMPVSSNTRRNEDGLGEPE GRASPDSPLTRWTKSLHSLLDGDQDGAYLFRFTFLEREKCVDTLDFWFACNGFRQMNLKD TKTLRVAKAIYKRYIENNSIVSKQLKPATKTYIRDGIKKQQIDSIMFDQAQTEIQSVM EENAYQMFLTSDIYLEYVRS GGENTAYMSNGGLGSLKVVCYGLPTLNEEEEWTCADFK CKLSPTVVLSSKTLRATASVRSTETVDSGYRSFKRSDPVNPNYHIGSGYVFAPATSAN DSEISSDALTDSDMSMTDSSVDGIPPYRVGSKKQLQREMHRSVKANGQVSLPHFPRTH RLPKEMTPVEPATFAAELISRLEKLELESRHSLEERLQQIREDEERESELTLSNR EGAPTQHPLSLLPSGSYEEDPQTILDDHLSRVLKTGCGSPGVGRYSPRSRSPDHHHH HHSQYHSLPPGGKLPAAASPGACPLLGGKGFVTKQTTKHVHHHYIHHHAVPKTKEE IEAEATQRVHCFCPGGSEYYCYCKSKSHSKAPETMPSEQFGGSRGSTLPKRNGKGTPE GLALPAREGGAP
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	100 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-Axin 2 Antibody (A14022)


Images:




Western blot analysis of extracts of various cell lines, using Anti-Axin 2 Antibody (A14022) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.



Immunohistochemistry analysis of paraffin-embedded rat brain using Anti-Axin 2 Antibody (A14022) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse spinal cord using Anti-Axin 2 Antibody (A14022) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of PC-12 cells using Anti-Axin 2 Antibody (A14022) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).