

Anti-MAP1B Antibody (A14425)

Specifications:

Name: Anti-MAP1B Antibody

Description: Rabbit polyclonal antibody to MAP1B.

Applications: WB, IHC, ICC/IF

Recommended Dilutions: WB: 1:200-1:2,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200

Reactivity: Human, Mouse, Rat

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids

1820-2100 of human MAP1B (NP_005900.2).

Sequence: PPIDAASAEPYGFRASVLFDTMQHHLALNRDLSTPGLEKDSGGKTPGDFSYAYQKPEE

TTRSPDEEDYDYESYEKTTRTSDVGGYYYEKIERTTKSPSDSGYSYETIGKTTKTPED GDYSYEIIEKTTRTPEEGGYSYDISEKTTSPPEVSGYSYEKTERSRRLLDDISNGYDD SEDGGHTLGDPSYSYETTEKITSFPESEGYSYETSTKTTRTPDTSTYCYETAEKITRT

PQASTYSYETSDLCYTAEKKSPSEARQDVDLCLVSSCEYKHPKTELSPS

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 350 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium

Azide.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




Anti-MAP1B Antibody (A14425)


Images:




Western blot analysis of extracts of various cell lines, using Anti-MAP1B Antibody (A14425) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Immunofluorescence analysis of C6 cells using Anti-MAP1B Antibody (A14425) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of NIH/3T3 cells using Anti-MAP1B Antibody (A14425) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of U2OS cells using Anti-MAP1B Antibody (A14425) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).