


Anti-ALDH3A1 Antibody (A14810)

Specifications:


Name:	Anti-ALDH3A1 Antibody
Description:	Rabbit polyclonal antibody to ALDH3A1.
Applications:	WB, ICC/IF
Recommended Dilutions:	WB: 1:500-1:1,000, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-300 of human ALDH3A1 (NP_001128640.1).
Sequence:	MSKISEAVKRARAAFSSGRTRPLQFRIQQLEALQRLIQEQEQELVGALAADLHKNAYY EEVVYVLEEIEYMIQKLPEWAADEPVEKTPQTQQDELYIHSEPLGVVLVIGTWNYPFN LTIQPMVGAIAAGNSVVLKPSELSENMASLLATIIPQYLDKDLYPVINGGVPETTELL KERFDHILYTGSTGVGKIIMTAAAKHLTPVTLELGGKSPCYVDKNCDLDVACRRIAWG KFMNSGQTCVAPDYILCDPSIQNQIVEKLLKSLKEFYGEDAKKSRDYGRIISARHFQR VMGLIEG
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	55 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-ALDH3A1 Antibody (A14810)


Images:




Western blot analysis of extracts of A-549 cells, using Anti-ALDH3A1 Antibody (A14810) at 1:700 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 5s.



Immunofluorescence analysis of rat stomach using Anti-ALDH3A1 Antibody (A14810) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of 293T cells using Anti-ALDH3A1 Antibody (A14810) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of A-549 cells using Anti-ALDH3A1 Antibody (A14810) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).