


Anti-TRBP Antibody (A15766)

Specifications:


Name:	Anti-TRBP Antibody
Description:	Rabbit polyclonal antibody to TRBP.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-366 of human TARBP2 (NP_599150.1).
Sequence:	MSEEEQSGSTTTGCGLPSIEQMLAANPGKTPISLLQEYGTTRIGKTPVYDLLKAEGQAH QPNFTFRVTVGDTSCGQGSPSKKAAKHKAAEVALKHLKGGSMLEPALEDSSSFPLDS SLPEDIPVFTAAAAATPVPSVVLTRSPPELQPPVSPQQSECNVPGALQELVVQKGWR LPEYTVTQESGPAHRKEFTMTCRVERFIEIGSGTSKKLAKRNAAKMLLRVHTVPLDA RDGNEVEPDDDHFSIGVGSRLDGLRNRGPGCTWDSLRSVGEKILSLRSCSLGSLGAL GPACCRVLSLSEEQAFHVSYLDIEELSLSGLCQCLVELSTQPATVCHGSATTREAR GEARRALQYLKIMAGSK
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	46 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-TRBP Antibody (A15766)


Images:




Western blot analysis of extracts of various cell lines, using Anti-TRBP Antibody (A15766) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 90s.



Immunohistochemistry analysis of paraffin-embedded human thyroid cancer using Anti-TRBP Antibody (A15766) at a dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human breast cancer tissue using Anti-TRBP Antibody (A15766) at a dilution of 1:200 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of A549 cells using Anti-TRBP Antibody (A15766).