

## **Anti-Cytochrome P450 1A2 Antibody (A17132)**

#### Specifications:

Name: Anti-Cytochrome P450 1A2 Antibody

Description: Rabbit polyclonal antibody to Cytochrome P450 1A2.

Applications: WB, ICC/IF

Recommended Dilutions: WB: 1:500-1:2,000, ICC/IF: 1:50-1:200

Reactivity: Human, Mouse, Rat

Immunogen: Recombinant fusion protein containing a sequence corresponding to amino acids 205-305

of human CYP1A2 (NP\_000752.2).

Sequence: FGQHFPESSDEMLSLVKNTHEFVETASSGNPLDFFPILRYLPNPALQRFKAFNQRFLW

FLQKTVQEHYQDFDKNSVRDITGALFKHSKKGPRASGNLIPQE

Host: Rabbit

Clonality: Polyclonal

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 54 kDa

Product Form: Liquid


Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium

Azide.

Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.

#### Images:




Western blot analysis of extracts of various cell lines, using Anti-Cytochrome P450 1A2 Antibody (A17132) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 90s.




# **Anti-Cytochrome P450 1A2 Antibody (A17132)**


### Images continued:



Immunofluorescence analysis of human liver cells using Anti-Cytochrome P450 1A2 Antibody (A17132) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of mouse liver cells using Anti-Cytochrome P450 1A2 Antibody (A17132) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of rat liver cells using Anti-Cytochrome P450 1A2 Antibody (A17132) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).