


Anti-Squamous Cell Carcinoma Antigen 1 Antibody [CPTC-SERPINB3-2] (A249946)

Specifications:


Name:	Anti-Squamous Cell Carcinoma Antigen 1 Antibody [CPTC-SERPINB3-2]
Description:	Mouse monoclonal [CPTC-SERPINB3-2] antibody to Squamous Cell Carcinoma Antigen 1.
Applications:	WB
Recommended Dilutions:	WB: 1-2 µg/ml
Reactivity:	Human
Immunogen:	Recombinant full-length human SERPINB3 protein.
Host:	Mouse
Clonality:	Monoclonal
Clone ID:	CPTC-SERPINB3-2
Isotype:	IgG1
Light Chains:	kappa
Conjugate:	Unconjugated
Purification:	Protein A/G chromatography.
Concentration:	200 µg/ml
Product Form:	Liquid
Formulation:	Supplied in 10mM Phosphate Buffered Saline with 0.05% BSA and 0.05% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
General Notes:	This monoclonal antibody is also available in a different formulation without BSA and Sodium Azide - Anti-Squamous Cell Carcinoma Antigen 1 Antibody [CPTC-SERPINB3-2] - BSA and Azide free (A253126).
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-Squamous Cell Carcinoma Antigen 1 Antibody [CPTC-SERPINB3-2] (A249946)

Images:



SDS-PAGE analysis of Anti-Squamous Cell Carcinoma Antigen 1 Antibody [CPTC-SERPINB3-2] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-Squamous Cell Carcinoma Antigen 1 Antibody [CPTC-SERPINB3-2]. Z-Score and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.