


Anti-ZAP70 Antibody [ZAP70/2035] (A250344)

Specifications:


Name:	Anti-ZAP70 Antibody [ZAP70/2035]
Description:	Mouse monoclonal [ZAP70/2035] antibody to ZAP70.
Specificity:	ZAP70 is a 70kDa protein tyrosine kinase found in T-cells and natural killer cells. Control of this protein translation is via the IgVH gene. ZAP70 protein is expressed in leukemic cells of approximately 25% of chronic lymphocytic leukemia (CLL) cases as well. Anti-ZAP70 expression is an excellent surrogate marker for the distinction between the Ig-mutated (anti-ZAP70 negative) and Ig-unmutated (anti-ZAP70 positive) CLL subtypes and can identify patient groups with divergent clinical courses. The anti-ZAP70 positive Ig-unmutated CLL cases have been shown to have a poorer prognosis.
Applications:	Flow Cytometry, IF, ELISA, IHC-P
Recommended Dilutions:	Flow Cytometry: 1-2 µg/million cells, IF: 1-2 µg/ml, IHC-P: 1-2 µg/ml
Reactivity:	Human
Immunogen:	Recombinant fragment, around amino acids 247-382, of human ZAP70 protein. The exact sequence is proprietary.
Host:	Mouse
Clonality:	Monoclonal
Clone ID:	ZAP70/2035
Isotype:	IgG1
Light Chains:	kappa
Conjugate:	Unconjugated
Purification:	Protein A/G chromatography.
Concentration:	200 µg/ml
Product Form:	Liquid
Formulation:	Supplied in 10mM Phosphate Buffered Saline with 0.05% BSA and 0.05% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
General Notes:	This monoclonal antibody is also available in a different formulation without BSA and Sodium Azide - Anti-ZAP70 Antibody [ZAP70/2035] - BSA and Azide free (A253524).
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-ZAP70 Antibody [ZAP70/2035] (A250344)


Images:




Immunohistochemical analysis of formalin-fixed, paraffin-embedded human tonsil using Anti-ZAP70 Antibody [ZAP70/2035].



SDS-PAGE analysis of Anti-ZAP70 Antibody [ZAP70/2035] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.



Immunofluorescent analysis of PFA fixed Jurkat cells stained with Anti-ZAP70 Antibody [ZAP70/2035] followed by Goat Anti-Mouse IgG (CF® 488) (Green). The nuclear counterstain is NucSpot (Red).



Flow cytometric analysis of PFA fixed Jurkat cells using Anti-ZAP70 Antibody [ZAP70/2035] followed by Goat Anti-Mouse IgG (CF® 488) (Blue). Isotype Control (Red).