


## Anti-TIGIT Antibody [TIGIT/3106] (Biotin) (A251215)

### Specifications:


Name:	Anti-TIGIT Antibody [TIGIT/3106] (Biotin)
Description:	Mouse monoclonal [TIGIT/3106] antibody to TIGIT (Biotin).
Specificity:	TIGIT is a checkpoint inhibitor which binds with high affinity to the poliovirus receptor (PVR), causing increased IL10 secretion, decreased IL12B secretion. TIGIT binding to PVR also causes the suppression of T cell activation by promoting the generation of mature immuno-regulatory dendritic cells. It is expressed at low levels on natural killer (NK) cells, as well as peripheral memory and regulatory CD4+ T cells. At the protein level, it is upregulated following the activation of these cells. Functionally, TIGIT is similar to CTLA4. The ligands for TIGIT include CD155 (signal abrogation) and CD226 (signal stimulation). It has been demonstrated to be upregulated on T cells in many cancers and is a immuno-oncology target for therapy.
Applications:	IHC-P
Recommended Dilutions:	IHC-P: 2-4 µg/ml
Reactivity:	Human
Immunogen:	Recombinant fragment, around amino acids 22-141, of human TIGIT protein. The exact sequence is proprietary.
Host:	Mouse
Clonality:	Monoclonal
Clone ID:	TIGIT/3106
Isotype:	IgG2b
Light Chains:	kappa
Conjugate:	Biotin
Purification:	Protein A/G chromatography.
Concentration:	100 µg/ml
Product Form:	Liquid
Formulation:	Supplied in 10mM Phosphate Buffered Saline with 0.05% BSA and 0.05% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

# Anti-TIGIT Antibody [TIGIT/3106] (Biotin) (A251215)


## Images:




Immunohistochemical analysis of formalin-fixed, paraffin-embedded human prostate carcinoma using Anti-TIGIT Antibody [TIGIT/3106] (Biotin).



Immunohistochemical analysis of formalin-fixed, paraffin-embedded human prostate carcinoma using Anti-TIGIT Antibody [TIGIT/3106] (Biotin).



SDS-PAGE analysis of Anti-TIGIT Antibody [TIGIT/3106] (Biotin) under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-TIGIT Antibody [TIGIT/3106] (Biotin). Z-Score and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.