


## Anti-ECM1 Antibody [rSPM217] - BSA and Azide free (A251572)

### Specifications:


Name:	Anti-ECM1 Antibody [rSPM217] - BSA and Azide free
Description:	Recombinant mouse monoclonal [rSPM217] antibody to ECM1.
Specificity:	This antibody reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelium, and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.
Applications:	Flow Cytometry, IF, IHC-P
Recommended Dilutions:	Flow Cytometry: 0.5-1 µg/million cells, IF: 1-2 µg/ml, IHC-P: 1-2 µg/ml
Reactivity:	Human, Rat
Immunogen:	Recombinant full-length human ECM1 protein.
Host:	Mouse
Clonality:	Monoclonal
Clone ID:	rSPM217
Isotype:	IgG1
Light Chains:	kappa
Conjugate:	Unconjugated
Purification:	Protein A/G chromatography.
Concentration:	1 mg/ml
Product Form:	Liquid
Formulation:	Supplied in 10mM Phosphate Buffered Saline; without Sodium Azide and carrier free.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
General Notes:	This monoclonal antibody is also available in a different formulation with BSA and Sodium Azide - Anti-ECM1 Antibody [rSPM217] (A248390).
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-ECM1 Antibody [rSPM217] - BSA and Azide free (A251572)

### Images:



Immunohistochemical analysis of formalin-fixed, paraffin-embedded human liver using Anti-ECM1 Antibody [rSPM217].



SDS-PAGE analysis of Anti-ECM1 Antibody [rSPM217] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.