


## Anti-FLI1 Antibody [FLI1/3183] - BSA and Azide free (A251733)

### Specifications:


Name:	Anti-FLI1 Antibody [FLI1/3183] - BSA and Azide free
Description:	Mouse monoclonal [FLI1/3183] antibody to FLI1.
Specificity:	This antibody recognizes a protein of 51kDa, which is identified as FLI1. This protein, a member of the ETS family of DNA binding transcription factors, is involved in cellular proliferation and tumorigenesis. Ets-1 is the prototype member of a family of genes identified on the basis of homology to the v-Ets oncogene isolated from the E26 erythroblastosis virus. Members of the Ets gene family share a highly conserved carboxy-terminal domain containing a sequence related to the SV40 large T antigen nuclear localization signal sequence. Approximately 90% of Ewings Sarcomas. High sensitivity and specificity of Fli-1 equals to or exceeds that of the established vascular markers like CD31, CD34, and Factor VIII.
Applications:	ELISA
Reactivity:	Human
Immunogen:	Recombinant fragment, around amino acids 107-261, of human FLI1 protein with hexa-histidine tag.
Host:	Mouse
Clonality:	Monoclonal
Clone ID:	FLI1/3183
Isotype:	IgG1
Light Chains:	kappa
Conjugate:	Unconjugated
Purification:	Protein A chromatography.
Concentration:	1 mg/ml
Product Form:	Liquid
Formulation:	Supplied in 10mM Phosphate Buffered Saline; without Sodium Azide and carrier free.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
General Notes:	This monoclonal antibody is also available in a different formulation with BSA and Sodium Azide - Anti-FLI1 Antibody [FLI1/3183] (A248551).
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

# Anti-FLI1 Antibody [FLI1/3183] - BSA and Azide free (A251733)

## Images:



SDS-PAGE analysis of Anti-FLI1 Antibody [FLI1/3183] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-FLI1 Antibody [FLI1/3183]. Z-Score and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.