

Anti-CD86 Antibody [C86/1146] - BSA and Azide free (A253825)

Specifications:

Name: Anti-CD86 Antibody [C86/1146] - BSA and Azide free

Description: Mouse monoclonal [C86/1146] antibody to CD86.

Specificity: This antibody recognizes a protein of 70kDa, which is identified as CD86. CD86 is a type I

transmembrane glycoprotein and a member of the immunoglobulin superfamily of cell surface receptors. It is expressed at high levels on resting peripheral monocytes and dendritic cells and at very low density on resting B and T lymphocytes. CD86 expression is rapidly upregulated by B cell specific stimuli with peak expression at 18 to 42 hours after stimulation. CD86, along with CD80/B71, is an important accessory molecule in T cell co-stimulation via its interaction with CD28 and CD152/CTLA4. Since CD86 has rapid kinetics of induction, it is the major CD28 ligand expressed early in the immune response. It is also found on malignant Hodgkin and Reed Sternberg (HRS) cells in Hodgkins disease.

Applications: IF, WB, IHC-P

Recommended Dilutions: IF: 1-2 μg/ml, WB: 1-2 μg/ml, IHC-P: 2-4 μg/ml

Reactivity: Human, Mouse, Rat

Immunogen: Recombinant full-length human CD86 protein.

Host: Mouse

Clonality: Monoclonal

Clone ID: C86/1146

Isotype: IgG1

Light Chains: kappa

Conjugate: Unconjugated

Purification: Protein A/G chromatography.

Concentration: 1 mg/ml

Product Form: Liquid

Formulation: Supplied in 10mM Phosphate Buffered Saline; without Sodium Azide and carrier free.

Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

General Notes: This monoclonal antibody is also available in a different formulation with BSA and Sodium

Azide - Anti-CD86 Antibody [C86/1146] (A250645).




Anti-CD86 Antibody [C86/1146] - BSA and Azide free (A253825)

Specifications continued:


Disclaimer:

This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Immunohistochemical analysis of formalin-fixed, paraffin-embedded human melanoma using Anti-CD86 Antibody [C86/1146].



Western blot analysis of Daudi cell lysate using Anti-CD86 Antibody [C86/1146].



Immunofluorescent analysis of Raji cells stained with Anti-CD86 Antibody [C86/1146] followed by Goat Anti-Mouse IgG (CF® 488) (Green). Nuclei are stained with RedDot.



SDS-PAGE analysis of Anti-CD86 Antibody [C86/1146] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.