

## Anti-CD35 Antibody [CR1/6377] - BSA and Azide free (A278151)

### Specifications:


|                        |  |
|------------------------|--|
| Name:                  | Anti-CD35 Antibody [CR1/6377] - BSA and Azide free   |
| Description:           | Mouse monoclonal [CR1/6377] antibody to CD35.  |
| Specificity:           | CD35, complement receptor 1, is a cell membrane-bound, monomeric glycoprotein on numerous cell types including erythrocytes, leukocytes, glomerular podocytes, and follicular dendritic cells. The primary function of CD35 is to serve as the cellular receptor for C3b and C4b, the most important components of the complement system leading to clearance of foreign macromolecules. CD35 antigen is found on erythrocytes, B cells, a subset of T cells, monocytes, as well as eosinophils, and neutrophils. Anti-CD35 is considered a mature B-cell marker which labels follicular dendritic reticulum cells and tumors derived from such cells such as follicular dendritic cell tumor/sarcoma. |
| Applications:          | IHC-P  |
| Recommended Dilutions: | IHC-P: 1-2 µg/ml   |
| Reactivity:            | Human  |
| Immunogen:             | Recombinant fragment, within amino acids 650-850, of human CD35 protein. The exact sequence is proprietary.  |
| Host:                  | Mouse  |
| Clonality:             | Monoclonal   |
| Clone ID:              | CR1/6377   |
| Isotype:               | IgG1   |
| Light Chains:          | kappa  |
| Conjugate:             | Unconjugated   |
| Purification:          | Protein A/G chromatography.  |
| Concentration:         | 1 mg/ml  |
| Product Form:          | Liquid   |
| Formulation:           | Supplied in 10mM Phosphate Buffered Saline; without Sodium Azide and carrier free.   |
| Storage:               | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.  |
| General Notes:         | This monoclonal antibody is also available in a different formulation with BSA and Sodium Azide - Anti-CD35 Antibody [CR1/6377] (A277563).   |

## Anti-CD35 Antibody [CR1/6377] - BSA and Azide free (A278151)


### Specifications continued:

**Disclaimer:** This product is for research use only. It is not intended for diagnostic or therapeutic use.


### Images:



SDS-PAGE analysis of Anti-CD35 Antibody [CR1/6377] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.



Immunohistochemical analysis of formalin-fixed, paraffin-embedded human spleen tissue using Anti-CD35 Antibody [CR1/6377] at 2µg/ml.



Immunohistochemical analysis of formalin-fixed, paraffin-embedded human tonsil tissue using Anti-CD35 Antibody [CR1/6377] at 2µg/ml.

## Anti-CD35 Antibody [CR1/6377] - BSA and Azide free (A278151)

Images continued:



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-CD35 Antibody [CR1/6377]. Z-Score and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.