

## Anti-ID1 Antibody [PCRP-ID1-2F11] - BSA and Azide free (A278240)

### Specifications:

Name:	Anti-ID1 Antibody [PCRP-ID1-2F11] - BSA and Azide free
Description:	Mouse monoclonal [PCRP-ID1-2F11] antibody to ID1.
Specificity:	Members of the Id family of basic helix-loop-helix (bHLH) proteins include Id1, Id2, Id3 and Id4. They are ubiquitously expressed and dimerize with members of the class A and B HLH proteins. Due to the absence of the basic region, the resulting heterodimers cannot bind DNA. The Id-type proteins thus appear to negatively regulate DNA binding of bHLH proteins. Since Id1 inhibits DNA binding of E12 and Myo D, it apparently functions to inhibit muscle-specific gene expression. Under conditions that facilitate muscle cell differentiation, the Id protein levels fall, allowing E12 and/or E47 to form heterodimers with Myo D and myogenin, which in turn activate myogenic differentiation. It has been shown that expression of each of the Id proteins is strongly dependent on growth factor activation and that reduction of Id mRNA levels by antisense oligonucleotides leads to a delayed reentry of arrested cells into the cell cycle following growth factor stimulation.
Applications:	ELISA, IP, Flow Cytometry, IF
Recommended Dilutions:	IP: 1-2 $\mu$ g / 100-500 $\mu$ g proteins, Flow Cytometry: 1-2 $\mu$ g/million cells, IF: 1-2 $\mu$ g/ml
Reactivity:	Human
Immunogen:	Recombinant fragment, around amino acids 5-150, of human ID1 protein. The exact sequence is proprietary.
Host:	Mouse
Clonality:	Monoclonal
Clone ID:	PCRP-ID1-2F11
Isotype:	IgG1
Conjugate:	Unconjugated
Purification:	Protein A/G chromatography.
Concentration:	1 mg/ml
Product Form:	Liquid
Formulation:	Supplied in 10mM Phosphate Buffered Saline; without Sodium Azide and carrier free.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.


# Anti-ID1 Antibody [PCRP-ID1-2F11] - BSA and Azide free (A278240)

## Specifications continued:


**General Notes:** This monoclonal antibody is also available in a different formulation with BSA and Sodium Azide - Anti-ID1 Antibody [PCRP-ID1-2F11] (A277652).

**Disclaimer:** This product is for research use only. It is not intended for diagnostic or therapeutic use.


## Images:



Flow cytometric analysis of PFA-fixed HeLa cells using Anti-ID1 Antibody [PCRP-ID1-2F11] followed by Goat Anti-Mouse IgG (CF® 488) (Blue). Isotype Control (Red).




Immunofluorescent analysis of PFA-fixed HeLa cells stained with Anti-ID1 Antibody [PCRP-ID1-2F11] followed by Goat Anti-Mouse IgG (CF® 488) (Green). CF® 640A Phalloidin (Red).



SDS-PAGE analysis of Anti-ID1 Antibody [PCRP-ID1-2F11] under non-reduced and reduced conditions; showing intact IgG and intact heavy and light chains, respectively. SDS-PAGE analysis confirms the integrity and purity of the antibody.

## Anti-ID1 Antibody [PCRP-ID1-2F11] - BSA and Azide free (A278240)

Images continued:



Analysis of protein array containing more than 19,000 full-length human proteins using Anti-ID1 Antibody [PCRP-ID1-2F11]. Z-Score and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target; a MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.