

## Anti-Fatty Acid Synthase Antibody [ARC0377] (A305468)

## Specifications:

Name: Anti-Fatty Acid Synthase Antibody [ARC0377]

Description: Rabbit monoclonal [ARC0377] antibody to Fatty Acid Synthase.

Applications: WB, IHC, ICC/IF, IP

Recommended Dilutions: WB: 1:500-1:2,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200, IP: 1:100-1:500

Reactivity: Human, Mouse

Immunogen: A synthetic peptide corresponding to a sequence within amino acids 2400-2500 of human

FASN (P49327).

Sequence: AVDLIIKSHQGLDRQELSFAARSFYYKLRAAEQYTPKAKYHGNVMLLRAKTGGAYGED

LGADYNLSQVCDGKVSVHVIEGDHRTLLEGSGLESIISIIHSS

Host: Rabbit

Clonality: Monoclonal

Clone ID: ARC0377

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 273 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02%

Sodium Azide.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




## **Anti-Fatty Acid Synthase Antibody [ARC0377] (A305468)**


## Images:




Western blot analysis of extracts of various cell lines, using Anti-Fatty Acid Synthase Antibody [ARC0377] (A305468) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 3min.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using Anti-Fatty Acid Synthase Antibody [ARC0377] (A305468) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of NIH-3T3 cells using Anti-Fatty Acid Synthase Antibody [ARC0377] (A305468) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of HeLa cells using Anti-Fatty Acid Synthase Antibody [ARC0377] (A305468) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).