

Anti-FBXO28 Antibody (A305504)


Specifications:

Name:	Anti-FBXO28 Antibody
Description:	Rabbit polyclonal antibody to FBXO28.
Applications:	WB
Recommended Dilutions:	WB: 1:500-1:2,000
Reactivity:	Human, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 91-312 of human FBXO28 (NP_055991.1).
Sequence:	CKRMDLVCQRMLNQGFLKVERYHNLCQKQVKAQLPRRESERRNHSLARHADILAAVET RLSLLNMTFMKYVDSNLCCFIPGKVIDEIYRVLRYVNSTRAPQRAHEVLQELRDISSM AMEYFDEKIVPILKRKLPGSDVSGRLMGSPPVPGPSAALTTMQLFSKQNPSRQEVTKL QQQVKTNGAGVTVLRREISELRTKVQEQQKQLQDQDQKLLEQTQIIGE
Host:	Rabbit
Host: Clonality:	Rabbit Polyclonal
Clonality:	Polyclonal
Clonality: Isotype:	Polyclonal IgG
Clonality: Isotype: Conjugate:	Polyclonal IgG Unconjugated
Clonality: Isotype: Conjugate: Purification:	Polyclonal IgG Unconjugated Affinity purification.
Clonality: Isotype: Conjugate: Purification: Molecular Weight:	Polyclonal IgG Unconjugated Affinity purification. 45 kDa
Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form:	Polyclonal IgG Unconjugated Affinity purification. 45 kDa Liquid
Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form: Formulation:	Polyclonal IgG Unconjugated Affinity purification. 45 kDa Liquid Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.

antibodies

Anti-FBXO28 Antibody (A305504)

Images:



Western blot analysis of extracts of various cell lines, using Anti-FBXO28 Antibody (A305504) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 60s.