antibodies

Anti-Fukutin Antibody [ARC2559] (A305661)


Specifications:

| Name: | Anti-Fukutin Antibody [ARC2559] |
|------------------------|--|
| Description: | Rabbit monoclonal [ARC2559] antibody to Fukutin. |
| · | |
| Applications: | WB, IHC |
| Recommended Dilutions: | WB: 1:500-1:1,000, IHC: 1:50-1:200 |
| Reactivity: | Human, Mouse, Rat |
| Immunogen: | A synthetic peptide corresponding to a sequence within amino acids 362-461 of human Fukutin (O75072). |
| Sequence: | DVKLDVFFFYEETDHMWNGGTQAKTGKKFKYLFPKFTLCWTEFVDMKVHVPCETLEYI EANYGKTWKIPVKTWDWKRSPPNVQPNGIWPISEWDEVIQLY |
| Host: | Rabbit |
| Clonality: | Monoclonal |
| Clone ID: | ARC2559 |
| lsotype: | lgG |
| Conjugate: | Unconjugated |
| Purification: | Affinity purification. |
| Molecular Weight: | 54 kDa |
| Product Form: | Liquid |
| Formulation: | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide. |
| Storage: | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |


antibodies

Anti-Fukutin Antibody [ARC2559] (A305661)

Images:



Western blot analysis of extracts of various cell lines, using Anti-Fukutin Antibody [ARC2559] (A305661) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded human colon tissue using Anti-Fukutin Antibody [ARC2559] (A305661) at a dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.