

Anti-NF-kB p65 (phospho Ser536) Antibody [ARC55585] (A305925)

Specifications:

Name: Anti-NF-kB p65 (phospho Ser536) Antibody [ARC55585]

Description: Rabbit monoclonal [ARC55585] antibody to NF-kB p65 (phospho Ser536).

Applications: WB

Recommended Dilutions: WB: 1:1,000-1:5,000

Reactivity: Human, Mouse, Rat

Immunogen: A synthetic phosphorylated peptide around S536 of human Phospho-NF-kB p65/RelA-S536

(NP_068810.3).

Sequence: FSSIA

Host: Rabbit

Clonality: Monoclonal

Clone ID: ARC55585

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 65 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.05%

Proclin 300.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




Anti-NF-kB p65 (phospho Ser536) Antibody [ARC55585] (A305925)

Images:



Western blot analysis of extracts of HeLa, using Anti-NF-kB p65 (phospho Ser536) Antibody [ARC55585] (A305925) at 1:2,000 dilution. HeLa cells were treated by TNF-alpha+50nM CA (by 100ng/ml TNF-alpha + 50nM CA) at 37°C for 30 minutes after serum-starvation overnightThe secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.



Western blot analysis of extracts of various lysates, using Anti-NF-kB p65 (phospho Ser536) Antibody [ARC55585] (A305925) at 1:2,000 dilution. NIH/3T3 and C6 were treated by TNF-alpha+50nM CA (100ng/ml) at 37°C for 30 minutes after serum-starvation overnight. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 60s.