

Anti-CDCA5 Antibody [ARC2096] (A306109)

Specifications:

Name: Anti-CDCA5 Antibody [ARC2096]

Description: Rabbit monoclonal [ARC2096] antibody to CDCA5.

Applications: WB, IHC

Recommended Dilutions: WB: 1:500-1:2,000, IHC: 1:50-1:200

Reactivity: Human

Immunogen: A synthetic peptide corresponding to a sequence within amino acids 153-252 of human

CDCA5 (Q96FF9).

Sequence: GSASTSTPGRRSCFGFEGLLGAEDLSGVSPVVCSKLTEVPRVCAKPWAPDMTLPGISP

PPEKQKRKKKMPEILKTELDEWAAAMNAEFEAAEQFDLLVE

Host: Rabbit

Clonality: Monoclonal

Clone ID: ARC2096

Isotype: IgG

Conjugate: Unconjugated

Purification: Affinity purification.

Molecular Weight: 35 kDa

Product Form: Liquid

Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02%

Sodium Azide.


Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.




Anti-CDCA5 Antibody [ARC2096] (A306109)

Images:



Western blot analysis of extracts of HepG2 cells, using Anti-CDCA5 Antibody [ARC2096] (A306109) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded human thyroid cancer using Anti-CDCA5 Antibody [ARC2096] (A306109) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.