


Anti-Von Willebrand Factor Antibody [ARC52173] (A306677)

Specifications:


Name:	Anti-Von Willebrand Factor Antibody [ARC52173]
Description:	Rabbit monoclonal [ARC52173] antibody to Von Willebrand Factor.
Applications:	IHC
Recommended Dilutions:	IHC: 1:500-1:1,000
Reactivity:	Human, Mouse
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 2645-2813 of human von Willebrand factor (VWF) (NP_000543.3).
Sequence:	LPTACTIQLRGGQIMTLKRDETLQDGCDFHCKVNERGEYFWEKRVTGCPPFDEHKCL AEGGKIMKIPGTCCDTCEEPECNDITARLQYVKVGSCKSEVEVDIHYCQGKCASKAMY SIDINDVQDQCSCCSPTRTEPMQVALHCTNGSVVYHEVLNAMECKCSPRKCSK
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC52173
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-Von Willebrand Factor Antibody [ARC52173] (A306677)


Images:




Immunohistochemistry analysis of paraffin-embedded human colon tissue using Anti-Von Willebrand Factor Antibody [ARC52173] (A306677) at a dilution of 1:800 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human tonsil using Anti-Von Willebrand Factor Antibody [ARC52173] (A306677) at a dilution of 1:800 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human esophageal cancer using Anti-Von Willebrand Factor Antibody [ARC52173] (A306677) at a dilution of 1:800 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse lung using Anti-Von Willebrand Factor Antibody [ARC52173] (A306677) at a dilution of 1:800 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.