

## Anti-FADD Antibody (A307154)


## Specifications:

Name:	Anti-FADD Antibody
Description:	Rabbit polyclonal antibody to FADD.
Applications:	WB, ICC/IF, IP
Recommended Dilutions:	WB: 1:500-1:1,000, ICC/IF: 1:50-1:200, IP: 1:500-1:1,000
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-205 of mouse FADD (NP_034305.1).
Sequence:	MDPFLVLLHSLSGSLSGNDLMELKFLCRERVSKRKLERVQSGLDLFTVLLEQNDLERG HTGLLRELLASLRRHDLLQRLDDFEAGTATAAPPGEADLQVAFDIVCDNVGRDWKRLA RELKVSEAKMDGIEEKYPRSLSERVRESLKVWKNAEKKNASVAGLVKALRTCRLNLVA DLVEEAQESVSKSENMSPVLRDSTVSSSETP
Host:	Rabbit
Host: Clonality:	Rabbit Polyclonal
Clonality:	Polyclonal
Clonality: Isotype:	Polyclonal IgG
Clonality: Isotype: Conjugate:	Polyclonal IgG Unconjugated
Clonality: Isotype: Conjugate: Purification:	Polyclonal IgG Unconjugated Affinity purification.
Clonality: Isotype: Conjugate: Purification: Molecular Weight:	Polyclonal IgG Unconjugated Affinity purification. 28 kDa
Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form:	Polyclonal IgG Unconjugated Affinity purification. 28 kDa Liquid


## antibodies

## Anti-FADD Antibody (A307154)


Images:




Western blot analysis of extracts of Mouse heart, using Anti-FADD Antibody (A307154) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.



Western blot analysis of extracts of Jurkat cells, using Anti-FADD Antibody (A307154) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.



Immunofluorescence analysis of mouse bone marrow cells using Anti-FADD Antibody (A307154) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunoprecipitation analysis of 600µg extracts of mouse heart cells using 3µg of Anti-FADD Antibody (A307154). This Western blot was performed on the immunoprecipitate using Anti-FADD Antibody (A307154) at a dilution of 1:1000.