


## Anti-Hexokinase II Antibody [ARC52057] (A307428)

### Specifications:


Name:	Anti-Hexokinase II Antibody [ARC52057]
Description:	Rabbit monoclonal [ARC52057] antibody to Hexokinase II.
Applications:	WB
Recommended Dilutions:	WB: 1:500-1:1,000
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-120 of human Hexokinase II (NP_000180.2).
Sequence:	MIASHLLAYFFTELNHDQVQKVDQYLYHMRLSDETLLEISKRFRKEMEKGLGATTHPT AAVKMLPTFVRSTPDGTEHGFLALDLGGTNFRVLWVKVTDNGLQKVEMENQIYAIP DIMR
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC52057
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	102 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.05% Proclin 300.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-Hexokinase II Antibody [ARC52057] (A307428)

### Images:



Western blot analysis of extracts from normal (control) and Hexokinase II knockout (KO) HeLa(KO) cells, using Anti-Hexokinase II Antibody [ARC52057] (A307428) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25  $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.



Western blot analysis of various lysates, using Anti-Hexokinase II Antibody [ARC52057] (A307428) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25  $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.