# antibodies

## Anti-PER2 Antibody [ARC1134] (A307556)


#### Specifications:

| Name:                  | Anti-PER2 Antibody [ARC1134]   |
|------------------------|--|
| Description:           | Rabbit monoclonal [ARC1134] antibody to PER2.  |
| Applications:          | WB, ICC/IF, IP, ChIP   |
| Recommended Dilutions: | WB: 1:500-1:2,000, ICC/IF: 1:50-1:200, ChIP: 1:50-1:200  |
| Reactivity:            | Human, Mouse   |
| Immunogen:             | A synthetic peptide corresponding to a sequence within amino acids 1-100 of human PER2 (O15055).         |
| Sequence:              | MNGYAEFPPSPSNPTKEPVEPQPSQVPLQEDVDMSSGSSGHETNENCSTGRDSQGSDC<br>DDSGKELGMLVEPPDARQSPDTFSLMMAKSEHNPSTSGCSSD |
| Host:                  | Rabbit   |
| Clonality:             | Monoclonal   |
| Clone ID:              | ARC1134  |
| lsotype:               | IgG  |
| Conjugate:             | Unconjugated   |
| Purification:          | Affinity purification.   |
| Molecular Weight:      | 140 kDa  |
| Product Form:          | Liquid   |
| Formulation:           | Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.     |
| Storage:               | Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.                    |
| Disclaimer:            | This product is for research use only. It is not intended for diagnostic or therapeutic use.             |


# antibodies

### Anti-PER2 Antibody [ARC1134] (A307556)


Images:



Western blot analysis of extracts of various cell lines, using Anti-PER2 Antibody [ARC1134] (A307556) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 3min.



Immunofluorescence analysis of HeLa cells using Anti-PER2 Antibody [ARC1134] (A307556) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of NIH-3T3 cells using Anti-PER2 Antibody [ARC1134] (A307556) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).