


Anti-Histone H4 Antibody [ARC2340] (A308423)

Specifications:


Name:	Anti-Histone H4 Antibody [ARC2340]
Description:	Rabbit monoclonal [ARC2340] antibody to Histone H4.
Applications:	WB, IHC
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 24-103 of human Histone H4 (P62805).
Sequence:	RDNIQGITKPAIRRLARRGGVKRISGLIYEETRGVLKVFLENVIRDAVITYTEHAKRKT VTAMDVVYALKRQGRTLYGFGG
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC2340
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	11 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-Histone H4 Antibody [ARC2340] (A308423)


Images:




Western blot analysis of extracts of various cell lines, using Anti-Histone H4 Antibody [ARC2340] (A308423) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded rat liver using Anti-Histone H4 Antibody [ARC2340] (A308423) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human colon tissue using Anti-Histone H4 Antibody [ARC2340] (A308423) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse kidney using Anti-Histone H4 Antibody [ARC2340] (A308423) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.