


Anti-GARS Antibody [ARC0514] (A308454)

Specifications:


Name:	Anti-GARS Antibody [ARC0514]
Description:	Rabbit monoclonal [ARC0514] antibody to GARS.
Applications:	WB
Recommended Dilutions:	WB: 1:500-1:1,000
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 640-739 of human GARS (P41250).
Sequence:	RHGVSHKVDSSSGSIGRRYARTDEIGVAFGVTIDFDTVNKTPHTATLRDRDSMRQIRA EISELPSIVQDLANGNITWADVEARYPLFEGQETGKKETIEE
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC0514
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	83 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-GARS Antibody [ARC0514] (A308454)

Images:



Western blot analysis of Jurkat, using Anti-GARS Antibody [ARC0514] (A308454) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.



Western blot analysis of extracts from wild type(WT) and GARS knockout (KO) 293T cells, using Anti-GARS Antibody [ARC0514] (A308454) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 180s.