


Anti-TAPP-1 Antibody [ARC2927] (A308627)

Specifications:


Name:	Anti-TAPP-1 Antibody [ARC2927]
Description:	Rabbit monoclonal [ARC2927] antibody to TAPP-1.
Applications:	WB
Recommended Dilutions:	WB: 1:100-1:500
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 300-404 of human TAPP1 (Q9HB21).
Sequence:	SEHPPGPSESKHAFRPTNAATATSHSTASRSNSLVSTFTMEKRGFYESLAKVKPGNFK VQTVSPREPASKVTEQALLRPQSKNGPQEKDCDLVLDLDDASLPVSDV
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC2927
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	45 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-TAPP-1 Antibody [ARC2927] (A308627)

Images:



Western blot analysis of extracts of various cell lines, using Anti-TAPP-1 Antibody [ARC2927] (A308627) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 90s.



Western blot analysis of extracts of various cell lines, using Anti-TAPP-1 Antibody [ARC2927] (A308627) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 120s.