


Anti-His Tag Antibody [DM1] - BSA and Azide free (A317439)

Specifications:

| | |
|------------------------|---|
| Name: | Anti-His Tag Antibody [DM1] - BSA and Azide free |
| Description: | Recombinant rabbit monoclonal [DM1] antibody to His Tag. |
| Applications: | ELISA, WB |
| Recommended Dilutions: | WB: 1:1,000 |
| Host: | Rabbit |
| Clonality: | Monoclonal |
| Clone ID: | DM1 |
| Isotype: | IgG |
| Conjugate: | Unconjugated |
| Purification: | Affinity chromatography. |
| Concentration: | Reconstitution dependent. |
| Product Form: | Lyophilized |
| Reconstitution: | Reconstitute with distilled sterile water. |
| Formulation: | Lyophilized from sterile Phosphate Buffered Saline, pH 7.4. Normally 5%-8% Trehalose is added as a protectant before lyophilization. |
| Storage: | Shipped at 4°C. Lyophilized: Store at -20°C to -80°C. Reconstituted: Aliquot and store at -80°C. Product is stable for one year. Avoid freeze/thaw cycles. |
| General Notes: | Prior to reconstitution, centrifuge the vial at 5,000g for 3-5 minutes at room temperature. Reconstitute with appropriate volume of distilled sterile water to bring product to 1mg/ml concentration. After addition of distilled sterile water, mix by gentle tapping. Note: It is not recommended to vortex or vigorously pipette the sample. |
| Disclaimer: | This product is for research use only. It is not intended for diagnostic or therapeutic use. |

Anti-His Tag Antibody [DM1] - BSA and Azide free (A317439)

Images:



Western blot analysis of His-Tag fusion protein using Anti-His Tag Antibody [DM1] - Azide free (A317439) at 1:1,000 dilution. Secondary Antibody: Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:5,000 dilution. His-tag fusion protein: 70kDa, 20ng per lane. Blocking buffer: 5% non-fat dry milk in TBST. Exposure time: 1s.