

Anti-Syndecan-1 Antibody [DM45] - BSA and Azide free (A318650)


Specifications:

Name:	Anti-Syndecan-1 Antibody [DM45] - BSA and Azide free
Description:	Recombinant rabbit monoclonal [DM45] antibody to Syndecan-1.
Applications:	ELISA, Flow Cytometry
Recommended Dilutions:	ELISA: 1:5,000-10,000, Flow Cytometry: 1:100
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	DM45
lsotype:	lgG
Conjugate:	Unconjugated
Purification:	Affinity chromatography.
Concentration:	Reconstitution dependent.
Product Form:	Lyophilized
Reconstitution:	Reconstitute with distilled sterile water.
Formulation:	Lyophilized from sterile Phosphate Buffered Saline, pH 7.4. Normally 5%-8% Trehalose is added as a protectant before lyophilization.
Storage:	Shipped at 4°C. Lyophilized: Store at -20°C to -80°C. Reconstituted: Aliquot and store at -80°C. Product is stable for one year. Avoid freeze/thaw cycles.
General Notes:	Prior to reconstitution, centrifuge the vial at 5,000g for 3-5 minutes at room temperature. Reconstitute with appropriate volume of distilled sterile water to bring product to 1mg/ml concentration. After addition of distilled sterile water, mix by gentle tapping. Note: It is not recommended to vortex or vigorously pipette the sample.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


antibodies

Anti-Syndecan-1 Antibody [DM45] - BSA and Azide free (A318650)


Images:



Expi 293 cell line was transfected with irrelevant protein (A) and human CD138 (B) were surface stained with Anti-Syndecan-1 Antibody [DM45] - BSA and Azide free (A318650) at 1µg/ml followed by Anti-Rabbit IgG Antibody (Alexa 488).



Flow cytometry analysis of antigen binding of Anti-Syndecan-1 Antibody [DM45] - BSA and Azide free (A318650). (A) Anti-Syndecan-1 Antibody [DM45] - BSA and Azide free (A318650) does not bind to Jurkat cells that do not express CD138. (B) A clear peak shift of Anti-Syndecan-1 Antibody [DM45] - BSA and Azide free (A318650) was seen compared to the control when incubated with CD138-expressing MM.1S cells, indicating strong binding of Anti-Syndecan-1 Antibody [DM45] - BSA and Azide free (A318650) to CD138. Antibodies were incubated at 5μg/ml.



Immunohistochemistry analysis of Anti-Syndecan-1 Antibody [DM45] - BSA and Azide free (A318650).