


Anti-SMYD1 Antibody (A80437)

Specifications:


Name:	Anti-SMYD1 Antibody
Description:	Rabbit polyclonal antibody to SMYD1.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:200-1:2,000, IHC: 1:20-1:200, ICC/IF: 1:20-1:100
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-350 of human SMYD1 (NP_938015.1).
Sequence:	MTIGRMENVEVFTAEGKGRGLKATKEFWAADIIFAERAYSAVVFDSLNVFVCHTCFKR QEKLHRCGQCKFAHYCDRTCQKDAWLNHKNECSAIKRYGKVPNENIRLAARIMWRVER EGTGLTEGCLSVDDLQNHVEHFGEEEQKDLRVDVDTFLQYWPPQSQQFSMQYISHIF GVINCNGFTLSDQRGLQAVGVGIFPNLGLVNHDCWPNCTVIFNNGNHEAVKSMFHTQM RIELRALGKISEGEELTVSYIDFLNVSEERKRQLKKQYYFDCTCEHCQKCLKDDLFLG VKDNPKPSQEVVKEMIQFSKDTLEKIDKARSEGLYHEVVKLCRECLEKQEPVFADTNI YM
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	57 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-SMYD1 Antibody (A80437)


Images:




Western blot analysis of extracts of various cell lines, using Anti-SMYD1 Antibody (A80437) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 60s.



Immunohistochemistry analysis of paraffin-embedded rat kidney using Anti-SMYD1 Antibody (A80437) at a dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human lung using Anti-SMYD1 Antibody (A80437) at a dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse heart using Anti-SMYD1 Antibody (A80437) at a dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.