


Anti-HDAC6 Antibody (A80508)

Specifications:


Name:	Anti-HDAC6 Antibody
Description:	Rabbit polyclonal antibody to HDAC6.
Applications:	WB, IHC, ICC/IF, IP
Recommended Dilutions:	WB: 1:100-1:500, IHC: 1:100-1:500, ICC/IF: 1:50-1:200, IP: 1:20-1:50
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 836-1104 of human HDAC6 (NP_006035.2).
Sequence:	VMKVEDREGPSSSKLVTKKAPQPAKPRLAERM TTTREKKVLEAGMGKVTSASFGEESTP GQTNSETAVVALTQDQPSEAATGGATLAQTISEAAIGGAMLGQTTSEEAVGGATPDQT TSEETVGGAILDQTTSEDAVGGATLGQTTSEEAVGGATLAQTTSEAAMEGATLDQTTS EEAPGGTELIQTPLASSTDHQTPTSPVQGTTPQISPSTLIGSLRTLLELGSESQGASE SQAPGEENLLGEAAGGQDMADSMMLMQGSRGLTDQAIF
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	160 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-HDAC6 Antibody (A80508)


Images:




Western blot analysis of extracts of Rat liver, using Anti-HDAC6 Antibody (A80508) at 1:500 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 μ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Immunohistochemistry analysis of paraffin-embedded human liver using Anti-HDAC6 Antibody (A80508) at a dilution of 1:400 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of HeLa cells using Anti-HDAC6 Antibody (A80508) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunoprecipitation analysis of 100 μ g extracts of HepG2 cells using 3 μ g of Anti-HDAC6 Antibody (A80508). This Western blot was performed on the immunoprecipitate using Anti-HDAC6 Antibody (A80508) at a dilution of 1:1000.