


Anti-Ionotropic Glutamate receptor 2 Antibody [ARC0572] (A81034)

Specifications:


Name:	Anti-Ionotropic Glutamate receptor 2 Antibody [ARC0572]
Description:	Rabbit monoclonal [ARC0572] antibody to Ionotropic Glutamate receptor 2.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	A synthetic peptide corresponding to a sequence within amino acids 150-250 of human GluR2/GRIA2 (P42262).
Sequence:	YLYDSDRGLSTLQAVLDSAAEKKWQVTAINVGNINNDKKDEMYRSLFQDLELKKERRV ILDCERDKVNDIVDQVITIGKHKVKGYHYIIANLGFTDGDLLKI
Host:	Rabbit
Clonality:	Monoclonal
Clone ID:	ARC0572
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	99 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol, 0.05% BSA, and 0.02% Sodium Azide.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

Anti-Ionotropic Glutamate receptor 2 Antibody [ARC0572] (A81034)


Images:




Western blot analysis of extracts of various cell lines, using Anti-Ionotropic Glutamate receptor 2 Antibody [ARC0572] (A81034) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 1s.



Immunohistochemistry analysis of paraffin-embedded rat brain using Anti-Ionotropic Glutamate receptor 2 Antibody [ARC0572] (A81034) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse brain using Anti-Ionotropic Glutamate receptor 2 Antibody [ARC0572] (A81034) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunofluorescence analysis of SH-SY5Y cells using Anti-Ionotropic Glutamate receptor 2 Antibody [ARC0572] (A81034) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).