

Anti-XPOT Antibody (A81114)


Specifications:

Description:Rabbit polyclonal antibody to XPOT.Applications:WBRecommended Dilutions:WB: 1:500-1:2,000Reactivity:Human, Mouse, RatImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human XPOT (NP_009166.2).Sequence:MDEQALLGLNPNADSDFRQRALAYFEQLKISPDAWQVCAEALAQRTYSDDHVKFFCFQ VLEHQVKYKYSELTTVQQQLIRETLISWLQAQMLNPQPEKTFIRNKAAQVFALLFVTE VLTKWPKFFFDILSVVDLNPRGVDLYLRILMAIDSELVDRDVVHTSEEARRNTLIKDT MREQCIPNLVESWYQILQNYQFTNSEHost:RabbitClonality:PolyclonalJotype:IgGPurification:Affnity purification.Molecular Weight:100 kDaInou KDaIuquidFornulation:Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.Storage:Shippet at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.	Name:	Anti-XPOT Antibody
Numerical stateWB: 1:500-1:2,000Reactivity:Human, Mouse, RatImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human XPOT (NP_009166.2).Sequence:MDEQALLGLNPNADSDFRQRALAYFEQLKISPDAWQVCAEALAQRTYSDDHVKFFCFQ VLEHQVKYKYSELTTVQQQLIRETLISWLQAQMLNPQPEKTFIRNKAAQVFALLFVTE YLTKWPKFFFDILSVVDLNPRGVDLYLRILMAIDSELVDRDVVHTSEEARRNTLIKDT MREQCIPNLVESWYQILQNYQFTNSEHost:RabbitClonality:PolyclonalIsotype:IgGConjugate:UnconjugatedPurification:Affnity purification.Molecular Weight:100 kDaIroduct Form:LiquidSupplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal	Description:	Rabbit polyclonal antibody to XPOT.
Reactivity:Human, Mouse, RatImmunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human XPOT (NP_009166.2).Sequence:MDEQALLGLNPNADSDFRQRALAYFEQLKISPDAWQVCAEALAQRTYSDDHVKFFCFQ VLEHQVKYKYSELTTVQQQLIRETLISWLQAQMLNPQPEKTFIRNKAAQVFALLFVTE YLTKWPKFFDILSVVDLNPRGVDLYLRILMAIDSELVDRDVVHTSEEARRNTLIKDT MREQCIPNLVESWYQILQNYQFTNSEHost:RabbitClonality:PolyclonalIsotype:IgGConjugate:UnconjugatedPurification:Affnity purification.Molecular Weight:10 kDaIroduct Form:LiquidKomulation:Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.	Applications:	WB
Immunogen:Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human XPOT (NP_009166.2).Sequence:MDEQALLGLNPNADSDFRQRALAYFEQLKISPDAWQVCAEALAQRTYSDDHVKFFCFQ VLEHQVKYKYSELTTVQQQLIRETLISWLQAQMLNPQPEKTFIRNKAAQVFALLFVTE YLTKWPKFFFDILSVVDLNPRGVDLYLRILMAIDSELVDRDVVHTSEEARRNTLIKDT MREQCIPNLVESWYQILQNYQFTNSEHost:RabbitClonality:PolyclonalIsotype:IgGConjugate:UnconjugatedPurification:Affinity purification.Molecular Weight:100 kDaProduct Form:LiquidSotypei CorrectSupplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.	Recommended Dilutions:	WB: 1:500-1:2,000
human XPOT (NP_009166.2).Sequence:MDEQALLGLNPNADSDFRQRALAYFEQLKISPDAWQVCAEALAQRTYSDDHVKFFCFQ VLEHQVKYKYSELTTVQQQLIRETLISWLQAQMLNPQPEKTFIRNKAAQVFALLFVTE YLTKWPKFFFDILSVVDLNPRGVDLYLRILMAIDSELVDRDVVHTSEEARRNTLIKDT MREQCIPNLVESWYQILQNYQFTNSEHost:RabbitClonality:PolyclonalIsotype:IgGConjugate:UnconjugatedPurification:Affinity purification.Molecular Weight:100 kDaProduct Form:LiquidSupplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.	Reactivity:	Human, Mouse, Rat
VLEHQVKYKYSELTTVQQQLIRETLISWLQAQMLNPQPEKTFIRNKAAQVFALLFVTE YLTKWPKFFFDILSVVDLNPRGVDLYLRILMAIDSELVDRDVVHTSEEARRNTLIKDT MREQCIPNLVESWYQILQNYQFTNSEHost:RabbitClonality:PolyclonalIsotype:IgGConjugate:UnconjugatedPurification:Affinity purification.Molecular Weight:100 kDaFormulation:Liquid	Immunogen:	
Clonality:PolyclonalIsotype:IgGConjugate:UnconjugatedPurification:Affinity purification.Molecular Weight:100 kDaProduct Form:LiquidSupple Ality purification purification.Formulation:Supple Ality purification purification.	Sequence:	VLEHQVKYKYSELTTVQQQLIRETLISWLQAQMLNPQPEKTFIRNKAAQVFALLFVTE YLTKWPKFFFDILSVVDLNPRGVDLYLRILMAIDSELVDRDVVHTSEEARRNTLIKDT
Isotype:IgGConjugate:UnconjugatedPurification:Affinity purification.Molecular Weight:100 kDaProduct Form:LiquidFormulation:Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.		
Conjugate:UnconjugatedPurification:Affinity purification.Molecular Weight:100 kDaProduct Form:LiquidFormulation:Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.	Host:	Rabbit
Purification:Affinity purification.Molecular Weight:100 kDaProduct Form:LiquidFormulation:Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.		
Molecular Weight:100 kDaProduct Form:LiquidFormulation:Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.	Clonality:	Polyclonal
Product Form: Liquid Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.	Clonality: Isotype:	Polyclonal IgG
Formulation: Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.	Clonality: Isotype: Conjugate:	Polyclonal IgG Unconjugated
	Clonality: Isotype: Conjugate: Purification:	Polyclonal IgG Unconjugated Affinity purification.
Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.	Clonality: Isotype: Conjugate: Purification: Molecular Weight:	Polyclonal IgG Unconjugated Affinity purification. 100 kDa
	Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form:	Polyclonal IgG Unconjugated Affinity purification. 100 kDa Liquid
Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.	Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form: Formulation:	Polyclonal IgG Unconjugated Affinity purification. 100 kDa Liquid Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Disclaiment	Clonality: Isotype: Conjugate: Purification: Molecular Weight:	Polyclonal IgG Unconjugated Affinity purification. 100 kDa

antibodies

Anti-XPOT Antibody (A81114)

Images:



Western blot analysis of extracts of various cell lines, using Anti-XPOT Antibody (A81114) at 1:3,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.