


Anti-FTCD Antibody (A82449)

Specifications:

Name:	Anti-FTCD Antibody
Description:	Goat polyclonal antibody to FTCD.
Applications:	ELISA, WB, IHC
Reactivity:	Human
Immunogen:	Synthetic peptide corresponding to Human FTCD (internal region).
Sequence:	CLREQGRGKDQPGRL
Host:	Goat
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Concentration:	100 µg at 0.5 mg/ml.
Product Form:	Liquid
Formulation:	Supplied in Tris Buffered Saline, pH 7.30, with 0.02% Sodium Azide and 0.5% BSA.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Anti-FTCD Antibody (A82449) (0.1µg/ml) staining of Human (A), (0.03µg/ml) of Mouse (B) and (0.3µg/ml) of Pig (C) Liver lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

Anti-FTCD Antibody (A82449)


Images continued:



Anti-FTCD Antibody (A82449) (3.75µg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Anti-FTCD Antibody (A82449) - Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10µg/ml) followed by Alexa Fluor 488 secondary antibody (2µg/ml), showing membrane and cytoplasmic and plasma membrane stain.



Anti-FTCD Antibody (A82449) - Immunofluorescence analysis of paraformaldehyde fixed HepG2 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10µg/ml) followed by Alexa Fluor 488 secondary antibody (2µg/ml), showing plasma membrane staining.