


### Anti-LDLRAP1 Antibody (A83944)

#### Specifications:

Name:	Anti-LDLRAP1 Antibody
Description:	Goat polyclonal antibody to LDLRAP1.
Applications:	ELISA, IF
Reactivity:	Human
Immunogen:	Synthetic peptide corresponding to Human LDLRAP1 (N terminal).
Sequence:	DALKSAGRALIRS-C
Host:	Goat
Clonality:	Polyclonal
lsotype:	lgG
Conjugate:	Unconjugated
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Concentration:	100 μg at 0.5 mg/ml.
Product Form:	Liquid
Formulation:	Supplied in Tris Buffered Saline, pH 7.30, with 0.02% Sodium Azide and 0.5% BSA.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

#### Images:




Anti-LDLRAP1 Antibody (A83944) Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 $\mu$ g/ml) followed by Alexa Fluor 488 secondary antibody (4 $\mu$ g/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 $\mu$ g/ml) followed by Alexa Fluor 488 secondary antibody (4 $\mu$ g/ml).

# antibodies

## Anti-LDLRAP1 Antibody (A83944)

Images continued:



Anti-LDLRAP1 Antibody (A83944) Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10µg/ml) followed by Alexa Fluor 488 secondary antibody (4µg/ml), showing showing cytoplasmic and some nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10µg/ml) followed by Alexa Fluor 488 secondary antibody (4µg/ml).