

Anti-FANCG Antibody (A83967)

Specifications:

Name: Anti-FANCG Antibody

Description: Goat polyclonal antibody to FANCG.

Applications: ELISA, WB, IHC

Reactivity: Human

Immunogen: Synthetic peptide corresponding to Human FANCG (C terminal).

Sequence: LEEFRTSLPKSCDL

Host: Goat

Clonality: Polyclonal

Isotype: IgG

Conjugate: Unconjugated

Purification: Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity

chromatography using the immunizing peptide.

Concentration: 100 µg at 0.5 mg/ml.

Product Form: Liquid

Formulation: Supplied in Tris Buffered Saline, pH 7.30, with 0.02% Sodium Azide and 0.5% BSA.

Storage: Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.

Disclaimer: This product is for research use only. It is not intended for diagnostic or therapeutic use.

Images:

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa Anti-FANCG Antibody (A83967) (0.5 μ g/ml) staining of HeLa cell lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

20kDa 15kDa




Anti-FANCG Antibody (A83967)


Images continued:



Anti-FANCG Antibody (A83967) ($1\mu g/ml$) staining of Jurkat nuclear lysate ($35\mu g$ protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



Anti-FANCG Antibody (A83967) ($3\mu g/ml$) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Anti-FANCG Antibody (A83967) (3 μ g/ml) staining of paraffin embedded Human Uterus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.