


Anti-HAVCR2 Antibody (A84174)

Specifications:

Name:	Anti-HAVCR2 Antibody
Description:	Goat polyclonal antibody to HAVCR2.
Applications:	ELISA, WB, IF, FC
Reactivity:	Human
Immunogen:	Synthetic peptide corresponding to Human HAVCR2 (internal region).
Sequence:	C-KWYSHSKEKIQN
Host:	Goat
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Concentration:	100 µg at 0.5 mg/ml.
Product Form:	Liquid
Formulation:	Supplied in Tris Buffered Saline, pH 7.30, with 0.02% Sodium Azide and 0.5% BSA.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.


Images:




Anti-HAVCR2 Antibody (A84174) (2µg/ml) staining of Jurkat (A), MOLT4 (B) and negative control A431 (C) cell lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.

Anti-HAVCR2 Antibody (A84174)


Images continued:



Anti-HAVCR2 Antibody (A84174) (1 μ g/ml) staining of Human Tonsil lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



Anti-HAVCR2 Antibody (A84174) - Immunofluorescence analysis of paraformaldehyde fixed HepG2 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (2 μ g/ml), showing membrane/cytoplasmic and nuclear staining.



Anti-HAVCR2 Antibody (A84174) - Flow cytometric analysis of paraformaldehyde fixed HepG2 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (1 μ g/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.