


### Anti-S100A10 Antibody (A88015)

#### Specifications:

Name:	Anti-S100A10 Antibody
Description:	Rabbit polyclonal antibody to S100A10.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:500-1:1,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-97 of human S100A10 (NP_002957.1).
Sequence:	MPSQMEHAMETMMFTFHKFAGDKGYLTKEDLRVLMEKEFPGFLENQKDPLAVDKIMKD LDQCRDGKVGFQSFFSLIAGLTIACNDYFVVHMKQKGKK
Host:	Rabbit
Clonality:	Polyclonal
lsotype:	lgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	11 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.05% Proclin 300.
Storage:	Shipped at $4^{\circ}$ C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

#### Images:




Western blot analysis of extracts of various cell lines, using Anti-S100A10 Antibody (A88015) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Enhanced Kit (RM00021). Exposure time: 180s.


# antibodies

## Anti-S100A10 Antibody (A88015)


Images continued:



Immunohistochemistry analysis of paraffin-embedded human esophagus using Anti-S100A10 Antibody (A88015) at a dilution of 1:20 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with IHC staining protocol.



Immunofluorescence analysis of PC-12 using Anti-S100A10 Antibody (A88015) at a dilution of 1:20 (40x lens). DAPI was used to stain the cell nuclei (blue).



Immunofluorescence analysis of A-431 using Anti-S100A10 Antibody (A88015) at a dilution of 1:20 (40x lens). DAPI was used to stain the cell nuclei (blue).