


## Anti-Spt6 Antibody (A88931)

### Specifications:


Name:	Anti-Spt6 Antibody
Description:	Rabbit polyclonal antibody to Spt6.
Applications:	WB, ICC/IF, IP
Recommended Dilutions:	WB: 1:500-1:2,000, ICC/IF: 1:50-1:200, IP: 1:100-1:500
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1500-1726 of human SUPT6H (NP_003161.2).
Sequence:	IFPTVNGLFRWFKDHYQDPVPGITPSSSSRTRTPASINATPANINLADLTRAVNALPQ NMTSQMFSAIAAVTGQGQNPATPAQWASSQYGYGGSSGGSSAYHVFPTPAQQPVATP LMTPSYSYTTSPQPITTPQYHQLQASTTPQSAQAQPQPSSSSRQRQQPKSNSHAAID WGKMAEQWLQEKEAERRKQKQRLTPRPSPSPMIESTPMSIAGDATPLLEMDR
Host:	Rabbit
Clonality:	Polyclonal
Isotype:	IgG
Conjugate:	Unconjugated
Purification:	Affinity purification.
Molecular Weight:	260 kDa
Product Form:	Liquid
Formulation:	Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.
Storage:	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
Disclaimer:	This product is for research use only. It is not intended for diagnostic or therapeutic use.

## Anti-Spt6 Antibody (A88931)


### Images:



Western blot analysis of extracts of various cell lines, using Anti-Spt6 Antibody (A88931) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25 $\mu$ g per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.



Immunofluorescence analysis of U-2 OS cells using Anti-Spt6 Antibody (A88931) at a dilution of 1:100. DAPI was used to stain the cell nuclei (blue).



Immunoprecipitation analysis of 600 $\mu$ g extracts of mouse heart using 3 $\mu$ g of Anti-Spt6 Antibody (A88931). This Western blot was performed on the immunoprecipitate using Anti-Spt6 Antibody (A88931) at a dilution of 1:500.