

## Anti-PREI3 Antibody (A88964)


### Specifications:

Name:	Anti-PREI3 Antibody
Description:	Rabbit polyclonal antibody to PREI3.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:500-1:2,000, IHC: 1:50-1:100, ICC/IF: 1:50-1:100
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1-225 of human MOB4 (NP_056202.2).
Sequence:	MVMAEGTAVLRRNRPGTKAQDFYNWPDESFDEMDSTLAVQQYIQQNIRADCSNIDKIL EPPEGQDEGVWKYEHLRQFCLELNGLAVKLQSECHPDTCTQMTATEQWIFLCAAHKTP KECPAIDYTRHTLDGAACLLNSNKYFPSRVSIKESSVAKLGSVCRRIYRIFSHAYFHH RQIFDEYENETFLCHRFTKFVMKYNLMSKDNLIVPILEEEVQNSVSGESEA
Host:	Rabbit
Host: Clonality:	Rabbit Polyclonal
Clonality:	Polyclonal
Clonality: Isotype:	Polyclonal IgG
Clonality: Isotype: Conjugate:	Polyclonal IgG Unconjugated
Clonality: Isotype: Conjugate: Purification:	Polyclonal IgG Unconjugated Affinity purification.
Clonality: Isotype: Conjugate: Purification: Molecular Weight:	Polyclonal IgG Unconjugated Affinity purification. 26 kDa
Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form:	Polyclonal IgG Unconjugated Affinity purification. 26 kDa Liquid


# antibodies

### Anti-PREI3 Antibody (A88964)


#### Images:



Western blot analysis of extracts of various cell lines, using Anti-PREI3 Antibody (A88964) at 1:3,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 5s.



Immunohistochemistry analysis of paraffin-embedded rat ovary using Anti-PREI3 Antibody (A88964) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.



Immunohistochemistry analysis of paraffin-embedded human oophoroma using Anti-PREI3 Antibody (A88964) at a dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.