

Anti-Robo1 Antibody (A88982)


Specifications:

Name:	Anti-Robo1 Antibody
Description:	Rabbit polyclonal antibody to Robo1.
Applications:	WB, IHC, ICC/IF
Recommended Dilutions:	WB: 1:200-1:2,000, IHC: 1:50-1:200, ICC/IF: 1:50-1:200
Reactivity:	Human, Mouse, Rat
Immunogen:	Recombinant fusion protein containing a sequence corresponding to amino acids 1452-1651 of human ROBO1 (NP_002932.1).
Sequence:	AAVMQKTRPAKKLKHQPGHLRRETYTDDLPPPPVPPPAIKSPTAQSKTQLEVRPVVVP KLPSMDARTDRSSDRKGSSYKGREVLDGRQVVDMRTNPGDPREAQEQQNDGKGRGNKA AKRDLPPAKTHLIQEDILPYCRPTFPTSNNPRDPSSSSSMSSRGSGSRQREQANVGRR NIAEMQVLGGYERGEDNNEELEETES
Host:	Rabbit
Host: Clonality:	Rabbit Polyclonal
Clonality:	Polyclonal
Clonality: Isotype:	Polyclonal IgG
Clonality: Isotype: Conjugate:	Polyclonal IgG Unconjugated
Clonality: Isotype: Conjugate: Purification:	Polyclonal IgG Unconjugated Affinity purification.
Clonality: Isotype: Conjugate: Purification: Molecular Weight:	Polyclonal IgG Unconjugated Affinity purification. 270 kDa
Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form:	Polyclonal IgG Unconjugated Affinity purification. 270 kDa Liquid
Clonality: Isotype: Conjugate: Purification: Molecular Weight: Product Form: Formulation:	Polyclonal IgG Unconjugated Affinity purification. 270 kDa Liquid Supplied in Phosphate Buffered Saline, pH 7.3, with 50% Glycerol and 0.01% Thiomersal.


antibodies

Anti-Robo1 Antibody (A88982)


Images:



Western blot analysis of extracts of various cell lines, using Anti-Robo1 Antibody (A88982) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 3s.



Western blot analysis of extracts of various cell lines, using Anti-Robo1 Antibody (A88982) at 1:1,000 dilution. The secondary antibody was Goat Anti-Rabbit IgG H&L Antibody (HRP) at 1:10,000 dilution. Lysates/proteins were present at 25µg per lane. The blocking buffer used was 3% non-fat dry milk in TBST. Detection was with a ECL Basic Kit. Exposure time: 10s.



Immunofluorescence analysis of U-2 OS cells using Anti-Robo1 Antibody (A88982) at a dilution of 1:100 (40x lens). DAPI was used to stain the cell nuclei (blue).